

Referenzen:

- Yoshihara A, Noh J, Yamaguchi T, Ohye H, Sato S, Sekiya K, Kosuga Y, Suzuki M, Matsumoto M, Kunii Y, Watanabe N, Mukasa K, Ito K, Ito K. Treatment of graves' disease with antithyroid drugs in the first trimester of pregnancy and the prevalence of congenital malformation. *J Clin Endocrinol Metab.* 2012 Jul;97(7):2396-403.
- Food and Drug Administration 2009: Information for Healthcare Professionals - Propylthiouracil-Induced Liver Failure (<http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/DrugSafetyInformationforHeathcareProfessionals/ucm162701.htm>.)
- Bahn RS, Burch HS, Cooper DS, Garber JR, Greenlee CM, Klein IL, Laurberg P, McDougall IR, Rivkees SA, Ross D, Sosa JA, Stan MN. The Role of Propylthiouracil in the Management of Graves' Disease in Adults: report of a meeting jointly sponsored by the American Thyroid Association and the Food and Drug Administration. *Thyroid.* 2009 Jul;19(7):673-4
- Food and Drug Administration 2010: Propylthiouracil tablets; Detailed View: Safety Labeling Changes Approved By FDA Center for Drug Evaluation and Research (CDER) – April 2010; BOXED WARNING (<http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm209256.htm>)
- Ono K, Kikuchi A, Takikawa KM, Hiroma T, Yoshizawa K, Sunagawa S, Takagi K, Momose Y, Nakamura T. Hernia of the umbilical cord and associated ileal prolapse through a patent omphalomesenteric duct: prenatal ultrasound and MRI findings. *Fetal Diagn Ther.* 2009;25(1):72-5.
- Foulds N, Walpole I, Elmslie F, Mansour S. Carbimazole embryopathy: an emerging phenotype. *Am J Med Genet A.* 2005 Jan 15;132A(2):130-5.
- Barbero P, Ricagni C, Mercado G, Bronberg R, Torrado M. Choanal atresia associated with prenatal methimazole exposure: three new patients. *Am J Med Genet A.* 2004 Aug 15;129A(1):83-6.
- Karg E, Bereg E, Gaspar L, Katona M, Turi S. Aplasia cutis congenita after methimazole exposure in utero. *Pediatr Dermatol.* 2004 Jul-Aug;21(4):491-4.
- Ferraris S, Valenzise M, Lerone M, Divizia MT, Rosaia L, Blaid D, Nemelka O, Ferrero GB, Silengo M. Malformations following methimazole exposure in utero: an open issue. *Birth Defects Res A Clin Mol Teratol.* 2003 Dec;67(12):989-92.
- Karlsson FA, Axelsson O, Melhus H. Severe embryopathy and exposure to methimazole in early pregnancy. *J Clin Endocrinol Metab.* 2002 Feb;87(2):947-9.
- Clementi M, Di Gianantonio E, Pelo E, Mammi I, Basile RT, Tenconi R. Methimazole embryopathy: delineation of the phenotype. *Am J Med Genet.* 1999 Mar 5;83(1):43-6.
- Abe M, Syuto T, Yokoyama Y, Ishikawa O. Aplasia cutis congenita after methimazole exposure in utero successfully treated with basic fibroblast growth factor. *Int J Dermatol.* 2010 Mar;49(3):334-5.
- Kannan L, Mishra S, Agarwal R, Kartikeyan V, Gupta N, Kabra M. Carbimazole embryopathy-bilateral choanal atresia and patent vitello-intestinal duct: a case report and review of literature. *Birth Defects Res A Clin Mol Teratol.* 2008 Sep;82(9):649-51.
- Clementi M, Di Gianantonio E, Cassina M, Leoncini E, Botto LD, Mastroiacovo P; SAFE-Med Study Group. Treatment of hyperthyroidism in pregnancy and birth defects. *J Clin Endocrinol Metab.* 2010 Nov;95(11):E337-41.
- Bahn RS, Burch HB, Cooper DS, Garber JR, Greenlee MC, Klein I, Laurberg P, McDougall IR, Montori VM, Rivkees SA, Ross DS, Sosa JA, Stan MN; American Thyroid Association; American Association of Clinical Endocrinologists. Hyperthyroidism and other causes of thyrotoxicosis: management guidelines of the American Thyroid Association and American Association of Clinical Endocrinologists. *Endocr Pract.* 2011 May-Jun;17(3):456-520.